




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 487/22	A1	(11) International Publication Number: WO 99/67249 (43) International Publication Date: 29 December 1999 (29.12.99)
<p>(21) International Application Number: PCT/US99/04924</p> <p>(22) International Filing Date: 5 March 1999 (05.03.99)</p> <p>(30) Priority Data: 09/102,394 22 June 1998 (22.06.98) US</p> <p>(71) Applicant: HEALTH RESEARCH, INC. [US/US]; Roswell Park Memorial Institute Division, Elm and Carlton Streets, Buffalo, NY 14263 (US).</p> <p>(72) Inventors: PANDEY, Ravindra, K.; 75 Lemay Court, Williamsville, NY 14221 (US). POTTER, William, R.; 2413 West River Road, Grand Island, NY 14072 (US). DOUGHERTY, Thomas, J.; 2306 West Oakfield, Grand Island, NY 14072 (US).</p> <p>(74) Agent: DUNN, Michael, L.; Dunn & Associates, P.O. Box 96, Newfane, NY 14108 (US).</p>	<p>(81) Designated States: CA, JP, MX, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> <p>Published With international search report.</p>	

(54) Title: ALKYL ETHER ANALOGS OF CHLORINS HAVING AN N-SUBSTITUTED IMIDE RING

(57) Abstract

Compounds having generic formula (I), where R¹, R² and R³ are independently alkyl of 3 through about 10 carbon atoms; provided that, R¹ and R² together contain at least six carbon atoms. The compounds have utility in photodynamic therapy in treatment of tumors and other diseases. The invention includes a method of treatment by contacting a tumor with the compound and then exposing the tumor to light.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

ALKYL ETHER ANALOGS OF CHLORINS HAVING AN N-SUBSTITUTED IMIDE RING

Background of the Invention

Some porphyrins and related tetrapyrrolic compounds, e.g. chlorins and bacteriochlorins, tend to be retained in malignant tumors in higher concentrations than in normal tissues. When the tetrapyrrolic compound is exposed to light of an appropriate wavelength, an excited state may occur and a singlet oxygen atom may be released. Such compounds which become excited by light are referred to herein as photodynamic compounds. When a patient is injected with an appropriate dose of a photodynamic compound, as previously described, the compound will concentrate in tumors which are present. The tumor may then be exposed to the appropriate wavelength of light to activate the compound which results in a tumoricidal effect. The release of singlet oxygen may be the cytotoxic species which, along with various other oxygen containing radicals, kills the tumor.

Porphyrins, chlorins, and bacteriochlorins including their analogs and derivatives have therefore recently found superior utility as photodynamic compounds for use in diagnosis and treatment of disease, especially certain cancers. These compounds have also found utility in treatment of psoriasis and papillomatosis.

Photofrin® (a porphyrin derivative), has been studied in most detail and is currently being used all over the world for treatment of various types of cancers. Despite the fact that it has already been approved for commercialization in Canada, Japan, and the United States, Photofrin photosensitizing agent has some disadvantages. In particular, it both photosensitizes the skin and lacks rapid normal tissue clearance, so patients must avoid exposure to sunlight for an extended period after its use. Further, it is a complex mixture of ether and ester linked dimers and higher oligomers, making it difficult to study mechanistically.

Photofrin photosensitizing agent also absorbs at a short wavelength making deep tissue penetration difficult and sometimes not possible. It is well established that both absorption and scattering of light by tissue increases as the wavelength decreases, and the most efficient photosensitizers are those which have strong absorption bands at the red end of the visible spectrum and the near IR region. Heme proteins accounts for most of the absorption of light in the visible region. Since this drops off rapidly beyond 550 nm, the effective depth of penetration doubles from 550 to 630 nm (where Photofrin® is activated) and doubles again in going to 700 nm, followed by 10% increase in penetration by moving into the near IR region (800 nm). However, further red shift is neither desirable (because of the lack of benefit from deeper penetration) nor easily achievable. Another reason to set the ideal wavelength in the range of 700-800 nm is due to the availability of diode lasers. Advantages of diode lasers are the low cost, negligible running cost, high reliability, small size and portability. Besides these considerations, tumor localization, dark toxicity,

phototoxicity, stability and formulation are also among the basic criteria to be considered for developing an efficient photosensitizer.

There is therefore still a quest for additional and new compounds for use in the field which have lower phototoxicity, better tissue clearance, better effectiveness and which absorb at higher wave lengths for better penetration.

Brief Description of the Drawings


FIGURE 1 is a schematic equation showing a method for preparation of compounds of the invention.

FIGURE 2 is a curve showing uptake in tumor versus skin fold of a purpurin-imide of the invention where R¹ is heptyl, R² is hexyl and R³ is methyl. The uptake is measured by optical density at the desired wavelength at 24 hours post injection.

Brief Description of the Invention

In accordance with the invention, a new series of compounds is therefore provided, which have good antitumor activity and which absorb at relatively long wave lengths of light.

More particularly, the compounds have the generic formula:



Where R^1 , R^2 , and R^3 are independently alkyl of 3 through about 10 carbon atoms; provided that, R^1 and R^2 together contain at least six carbon atoms. R^3 is preferably methyl or ethyl and R^2 and R^3 are preferably alkyl of 3 through 8 carbon atoms.

Detailed Description of the Invention

In accordance with the invention, a compound is provided having the generic structure described above.

Among the alkyl ether derivatives of pyropheophorbide-a, it has been shown that the biological activity increased by increasing the length of the carbon chain, being maximum with n-hexyl and n-heptyl analogs, further increase in the length of carbon chain reduced the in vivo efficacy.

In order to find the effect of such substituents in other series of compounds, in accordance with the invention, a series of various alkyl ether analogs of chlorins with N-substituted cyclic imides (purpurin-18 imides) were prepared. These compounds have long wavelength absorption at 705 nm, and offer considerable latitude in testing the modeling methods as well as refining it since they offer the possibility of two sites for alkyl substitution (R^1 and R^2) and allow the log P value to be held constant while changing the size and shape of the molecules. For example, the calculated (PALLAS) log P for chlorin where R^1 =propyl and R^2 =decyl and vice versa and where R^1 =heptyl and R^2 =hexyl and vice versa are identical (10.72) (i.e. each with 11 methylene groups and 2 terminal methyl groups). Numerous other examples are obvious. Thus, we can hold constant or vary log P while changing size and steric factors as well.

A method for preparation of the compounds of the invention is shown in the schematic equations set out in Figure 1.

R¹ is alkyl of 3 through about 10 and preferably 3 to 8 carbon atoms. R² is alkyl of 3 through about 10 carbon atoms and preferably 3 to 8 carbon atoms. R³ is methyl or ethyl. R¹ and R² together contain at least 6 carbon atoms. The alkyl groups may be linear or branched chain and may be substituted with other groups, e.g., alkoxy, halogen, ester or aromatic substituents.

Methylpheophorbide-a was isolated from the alga *Spirulina Pacifica*, and reacted with alkyl amines. The intermediate amides as carboxylic acid analogs were converted to the corresponding methyl esters, which on stirring with methanolic KOH at room temperature (5-10 min) produced the N-alkyl imide derivatives in 65-70% yield. The vinyl groups were then converted to various alkyl ether analogs by first reacting with 30% HBr/AcOH, and then with the desired alcohol (for example 4 to 6, Figure 1). The desired compounds (Table 1) were prepared in high yield (70-75%). The methyl ester group (e.g., heptyl ether analog of N-hexyl purpurin imide 4) can be hydrolyzed to the corresponding carboxylic acid.


Various compounds of the present invention were compared for in vivo tumor response, as shown in Table 1.

Radiation induced fibrosarcoma (RIF) tumors, were implanted subcutaneously into the axilla of 5-7 week old female C3H mice. When tumors grew to about 5mm in diameter, the mice were injected with various doses of the photosensitizers. The mice (6 mice per group) were restrained in aluminum holders and each tumor illuminated with 135 J/cm² of light from a laser tuned at the longest

wavelength absorption maximum of the photosensitizers at 705nm. The percentage of short term control was recorded daily.

The results are set forth in Table 1. The data indicated that: (1) the hydrophobicity of the molecule can be varied by changing the length of the carbon chain (N-alkyl or -alkyl substituents), (ii) in some cases, compounds with same hydrophobic characteristics (same log p values), did not show similar activity. (iii) thus besides hydrophobicity, the steric and electronic factors also play important role in designing effective photosensitizers.

In this series, the heptyl ether derivative of purpurin imide bearing N-hexyl substituent was found to be most effective at a dose of 1.0 $\mu\text{mol/kg}$.


Table 1. Preliminary *in vivo* Activity of Alkyl Ether Analogs of Purpurin-18 Imides

R ¹	R ²	Partition Coefficient (log P)	Tumor Response % [days]*						
			1-2	7	14	21	30	90	
Methyl	Propyl	6.22							NR
Propyl	Propyl	7.22	100	33	17				ongoing
Methyl	Hexyl	7.72							NR
Propyl	Hexyl (0.5 μmol)*	8.72							NR
Hexyl	Propyl (0.5 μmol)*	8.72	100	66	33	17	17	17	
Hexyl	Propyl	8.72	100	83	83	50	33	33	
Heptyl	Hexyl (0.25 μmol)*	10.72	100	60	20	20	20	20	
Heptyl	Hexyl	10.72	100	100	100	100	100	100	83

*Six mice/group (RIF tumor). 1.0 μmol/kg. 705 nm, 135J/cm² (24h post injection)

WHAT IS CLAIMED IS:

1. A compound of the formula:



Where R¹, R² and R³ are independently alkyl of 3 through about 10 carbon atoms; provided that, R¹ and R² together contain at least 6 carbon atoms.

2. The compound of Claim 1 wherein R¹ and R² are alkyl of 3 through 8 carbon atoms and R³ is methyl or ethyl.
3. The compound of Claim 1 wherein R¹ is heptyl, R² is hexyl and R³ is methyl.
4. A method for the treatment of a tumor which comprises contacting it with a compound of Claim 1 and exposing it to light having a wavelength of from about 650 to about 750 nm.

5. A method for the treatment of a tumor which comprises contacting it with a compound of Claim 2 and exposing it to light having a wavelength of from about 650 to about 750 nm.
6. A method for the treatment of a tumor which comprises contacting it with a compound of Claim 3 and exposing it to light having a wavelength of from about 650 to about 750 nm.
7. The method of Claim 4 wherein the light has a wavelength of from about 695 to about 710 nm.
8. The method of Claim 5 wherein the light has a wavelength of from about 695 to about 710 nm.
9. The method of Claim 6 wherein the light has a wavelength of from about 695 to about 710 nm.




FIGURE-1

Purpurin-Imide




FIGURE 2

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/04924

A. CLASSIFICATION OF SUBJECT MATTER IPC(6) :C07D 487/22 US CL :540/145, 472, 474 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 540/145, 472, 474 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,591,847 A (PANDEY et al) 07 January 1997, whole document	1-9
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: *A* document defining the general state of the art which is not considered to be of particular relevance *B* earlier document published on or after the international filing date *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) *O* document referring to an oral disclosure, use, exhibition or other means *P* document published prior to the international filing date but later than the priority date claimed *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art *Z* document member of the same patent family		
Date of the actual completion of the international search 09 APRIL 1999		Date of mailing of the international search report 27 APR 1999
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer PK SRIPADA Telephone No. (703) 308-1235